



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Enthalpy Diffusion in Multicomponent Flows

A. W. Cook

January 23, 2009

APS/DFD

Minneapolis, MN, United States

November 22, 2009 through November 24, 2009

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

Enthalpy Diffusion in Multicomponent Flows



Wake Vortex Study at Wallops Island



The Siege and Destruction of Jerusalem by the Romans Under the Command of Titus, A.D. 70, by David Roberts (1850)

Andrew W. Cook

APS/DFD, Minneapolis, Nov. 22-24, 2009

This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.

UCRL-PRES-??????

Do the multicomponent Euler equations adequately describe turbulent mixing?



$$\frac{\partial \rho Y_i}{\partial t} + \nabla \cdot (\rho Y_i \mathbf{u}) = 0$$

$$\frac{\partial \rho \mathbf{u}}{\partial t} + \nabla \cdot (\rho \mathbf{u} \mathbf{u} + p \underline{\underline{\mathbf{a}}}) = 0$$

$$\frac{\partial \rho E}{\partial t} + \nabla \cdot [(E + p) \mathbf{u}] = 0$$



Can these equations accurately predict temperature?



Can the multicomponent Navier-Stokes equations accurately predict temperature?



$$\frac{\partial \rho Y_i}{\partial t} + \nabla \cdot (\rho Y_i \mathbf{u}) = -\nabla \cdot \mathbf{J}_i$$

$$\frac{\partial \rho \mathbf{u}}{\partial t} + \nabla \cdot (\rho \mathbf{u} \mathbf{u} + p \underline{\underline{\mathbf{a}}}) = \nabla \cdot \underline{\underline{\hat{\mathbf{o}}}}$$

$$\frac{\partial \rho E}{\partial t} + \nabla \cdot [(E + p) \mathbf{u}] = \nabla \cdot (\underline{\underline{\hat{\mathbf{o}}}} \cdot \mathbf{u} - \mathbf{q}_c - \mathbf{q}_d)$$




Enthalpy Diffusion: $\mathbf{q}_d = h_i \mathbf{J}_i$

The species diffusion flux (J_i) is present in any simulation wherein mixing occurs.




J_i can represent:




- molecular diffusion (DNS, physical diffusivity)
- numerical diffusion (Euler solvers, ILES)
- subgrid-scale diffusion (LES, grid-scale transfer)
- turbulent diffusion (RANS, k - ϵ & k - l models)

q_d must balance J_i

The role of q_d is illustrated through a simple gedanken experiment.





If $q_d=0$ there can be no net mass flux; hence, $n_L M_L = n_H M_H$




$$\Delta S = \Delta S^< + \Delta S^> < 0$$


Shear layers evolve differently, depending on the presence of q_d .




Temperature can be extremely sensitive to enthalpy diffusion.




Enthalpy diffusion generates acoustic noise.



The Haas-Sturtevant shock-bubble experiment provides a good test of the importance of q_d .



Temperature can be extremely sensitive to enthalpy diffusion.



W.S. Don and C.B. Quillen, Numerical simulation of shock-cylinder interactions
J. Comput. Phys., 122:244-265 (1995)...

“Haas and Sturtevant's shock-helium cylinder interaction is well simulated by an Euler code”

“the temperature T is found to require a slightly heavier smoothing”

For Hydrogen-Air case...

“the general features are quite similar for both the Euler and the reactive Navier-Stokes simulations”

but

T (ambient) = 1000 K


T (ignition) = 853 K

Enthalpy diffusion indirectly results in a smoother density field.




OFF

ON




Density does not diffuse! Smoothing of the density field is brought about by a local divergence in the velocity field, which is here influenced by both heat conduction and enthalpy diffusion.

The Rayleigh-Taylor instability evolves differently, depending on the presence of q_d .




$$A = 0.87$$

Including q_d , $\Delta T < 0.0052$ eV
Excluding q_d , $0.5 \text{ eV} < T < 2 \text{ eV}$



$T_o = 1 \text{ eV}$ (to match Dimonte-Tipton, PF 18:085101)

Enthalpy diffusion generates acoustic noise in R-T instability.



Conclusions

Rudolf Clausius
Originator of the
concept of entropy



1. Enthalpy diffusion preserves the second law.
2. Euler solvers will not produce correct temperatures in mixing regions.
3. Navier-Stokes solvers will only produce correct temperatures if q_d is included.
4. Errors from neglecting enthalpy diffusion are most severe when differences in molecular weights are large.
5. In addition to temperature, enthalpy diffusion affects density, dilatation and other fields in subtle ways.
6. Reacting flow simulations that neglect the term are a dubious proposition.
7. Turbulence models for RANS and LES closures should preserve consistency between energy and species diffusion.